

Fate of transgenic DNA and evaluation of metabolic effects in goats fed genetically modified soybean and in their offsprings

R. Tudisco¹, V. Mastellone², M. I. Cutrignelli¹, P. Lombardi^{3†}, F. Bovera¹, N. Mirabella³, G. Piccolo¹, S. Calabrò¹, L. Avallone³ and F. Infascelli¹

¹Dipartimento di Scienze Zootecniche e Ispezione degli Alimenti, sez. B. Ferrara – Università di Napoli “Federico II”, Via F. Delpino, 1–80137 Napoli, Italy;

²Dipartimento di Medicina Sperimentale e Clinica “Gaetano Salvatore”, Università di Catanzaro Magna Græcia, via Tommaso Campanella, 115 – 88100 Catanzaro, Italy; ³Dipartimento di Strutture, Funzioni e Tecnologie Biologiche, Università di Napoli “Federico II”, Via F. Delpino, 1–80137 Napoli, Italy

(Received 16 March 2009; Accepted 29 March 2010)

The presence of DNA fragments in blood and milk from goats fed conventional (control) or Roundup Ready[®] soybean meal solvent extracted (s.e.; treated) was investigated by using a polymerase chain reaction approach. The same investigation was carried out on blood, skeletal muscle and organs from kids of both groups fed only dams' milk until weaning. Moreover, the possible effects on cell metabolism were evaluated by determination of several specific enzymes in serum, heart, skeletal muscle, liver and kidney. Fragments of the multicopy chloroplast (trnL) gene were found in blood and milk samples from goats of both groups. In kids, the chloroplast fragments were found in samples of both groups. In samples, which proved positive for the presence of chloroplast DNA, fragments of the specific soybean single copy gene (lectin) were detected in several blood and milk samples. The same fragment was also found in control and treated groups of kids. Transgenic fragments were not found in those samples, which were found positive for chloroplast fragments of control groups of either goats or kids. On the contrary, in blood and milk of treated goats, fragments both of the 35S promoter and the CP4 EPSPS gene were detected. These fragments were also found in treated kids with a significant detection of the 35S promoter in liver, kidney and blood, and of the CP4 EPSPS gene fragment in liver, kidney, heart and muscle. A significant increase in lactic dehydrogenase, mainly concerning the lactic dehydrogenase-1 isoenzyme was found in heart, skeletal muscle and kidney of treated kids, thus suggesting a change in the local production of the enzyme. Finally, no significant differences were detected concerning kid body and organ weight.

Keywords: genetically modified soybean meal, goat, kid, DNA passage

Implications

No direct evidence has been reported about animal health effects of genetically modified (GM) feed. Although the commercialisation of several GM feeds, including RoundUp Ready[®] soybean, has been authorised by the European authorities, concerns over safety still persist in the public, mainly regarding either the detection of transgenic plant genes and proteins in animal systems or allergenicity and toxicity of GM plants. The aims of this study were to evaluate the presence of plant DNA fragments and the levels of some health parameters in animals fed on GM soybean.

Introduction

In recent years, genetic engineering has been widely applied to agriculture in order to obtain specific plant characteristics,

which can lead to an improvement in both food quality and production. In this context, the European Food Safety Authority has stated that no recombinant DNA sequences have been found in any organ or tissue sample from animals fed genetically modified (GM) plants (EFSA, 2007). On this basis, a number of GM products have been approved for human consumption, but concerns over safety persist in the public, mainly regarding either the detection of transgenic plant genes and proteins in animal systems or allergenicity and toxicity of GM plants. Furthermore, some authors have shown the possible survival of recombinant plant DNA fragments after digestion, although health implications as well as biological relevance of data are still far from being confirmed (Nemeth *et al.*, 2004; Mazza *et al.*, 2005; Sharma *et al.*, 2006). The absorption of plant DNA across the intestinal barrier is a natural event, as demonstrated by the detection of endogenous plant genes in several animal tissues and products (Reuter and Aulrich, 2003; Tony *et al.*, 2003; Tudisco *et al.*, 2006b).

[†] E-mail: pilombar@unina.it

In a preliminary study (Laudadio *et al.*, 2006) concerning possible plant DNA fragment survival after digestive processes and their transfer and accumulation in tissues and organs of offspring, we found the chloroplast sequence, but no transgenic fragments in blood and tissues from lambs fed dam's milk until weaning. No relations between GM food and health have been reported to date.

Many feeding trials have been reported in which GM foods have been administered to rats or mice for prolonged periods, and parameters such as body weight (BW), feed consumption, blood chemistry, organ weights and histopathology have been measured. The majority of these experiments indicated no clinical effects nor histopathological abnormalities in organs or tissues of exposed animals (EFSA, 2008). By contrast, Malatesta *et al.* (2002) found significant nuclear modifications of hepatocytes in young and adult (2 to 8 months of age) female mice fed on GM soybean. In particular, GM-fed mice show irregularly shaped nuclei, which generally represents an index of high metabolic rate, and a higher number of nuclear pores, suggesting intense molecular trafficking. More recently, the same authors (Malatesta *et al.*, 2008) showed that several proteins belonging to hepatocyte metabolism, stress response, calcium signalling and mitochondria were expressed in 24-month-old female mice fed on GM soybean, indicating a more marked expression of senescence markers in comparison to controls. Moreover, hepatocytes of GM-fed mice showed mitochondrial and nuclear modifications indicative of reduced metabolic rates. Furthermore, Tudisco *et al.* (2006a) hypothesised that cell metabolism of several enzymes was altered in rabbit fed on GM soybean.

The aims of this study were to investigate the presence of plant DNA fragments in blood and milk from goats fed GM soybean and in tissues and organs from their kids fed only mother's milk for 60 days, and to study the activity of organ-specific enzymes.

Material and methods

Diets, animals and feeding

Twenty pregnant dairy goats were equally assigned to control (C) and treated (T) groups, homogeneous in parity and milk production at the previous lactation. Animals from different groups were housed in separate sheds. Experiments started 2 months before kidding; the animals were fed a diet consisting of oat hay and concentrate [crude protein (CP) 18% of dry matter (DM); metabolisable energy (MJ/DM) 12.22], the latter containing solvent extracted (s.e.) soybean meal (13% of concentrate DM) which was from conventional or GM (RoundUp Ready® (RR)) soy beans for groups C and T, respectively. RR is tolerant to the glyphosate family of herbicides by expressing transgenic DNA from the CP4 strain of *Agrobacterium tumefaciens* (CP4 EPSPS), encoding 5-enolpyruvylshikimate-3-phosphate synthase protein (glyphosate-tolerant soybean GTS 40-3-2; Padgett *et al.*, 1995). A polymerase chain reaction (PCR)-end point reaction confirmed the presence of p35S and CP4 EPSPS specific transgenes in the treated diet as well as their absence in the

Table 1 Chemical composition (g/kg DM) and ME (MJ/kg DM) of hays and concentrate

	Oat hay	Alfalfa hay	Concentrate*
CP	55.0	160.0	180
Ether extract	19.0	20.0	30.0
NDF	682.0	493.0	270.0
ADF	451.0	341.0	115.0
ADL	61.0	50.0	30.0
ME	6.36	9.39	12.22

DM = dry matter; ME = metabolisable energy.

*Ingredients (g/Kg DM): soft wheat bran 30; soybean s.e. 13; corn meal 13; sunflower meal 10.5; citrus pulp 8; sugar beet pulp 7.9; corn gluten feed 7; sugarcane molasses 7.5; CaCO₃ 1.5; CaHPO₄ 0.7; vitamins 0.2; NaCl 0.7.

control diet. Both groups received oat hay *ad libitum*, while the concentrate was administered in amounts of 200, 300 and 400 g/head per day, 60, 30 and 15 days before kidding, respectively. After kidding, administration of concentrate was gradually increased up to 700 g/head per day and oat hay was replaced by alfalfa hay in order to increase diet protein content. Water was given *ad libitum*. The chemical composition (Van Soest *et al.*, 1991; AOAC, 2000) and metabolisable energy (INRA, 1978) of hays and concentrate as well as ingredients of concentrate are reported in Table 1.

All goats had twin deliveries each of them with at least one male kid. Immediately after kidding, 10 male kids, were randomly selected from each group and allowed into individual cages positioned in a separate room. Each kid was fed only milk from its mother using a milk feeder. Milk was collected from each goat into sterile tubes with aseptic techniques and administered twice/day until 12 h before slaughtering. Kids were slaughtered at 60 ± 7 days of age (11.2 ± 1.1 kg live weight). Kid BWs were taken at birth and immediately before slaughtering.

Sample collection

In all, 20 dams (ten per group) with their kids were included in the trial. After kidding, 100 ml of milk from each goat (obtained by mixing the production of the two daily milkings) were collected at days 15, 30, 45, 60 and 75. Samples were immediately frozen on dry ice before storage at -20°C. Blood was withdrawn every month (day -60 until day 60) from each goat via jugular vein puncture. Blood was removed from kids before slaughter. In both cases, samples used for DNA extraction were added to tubes containing K3-EDTA (ethylene diamine tetra acetic acid) 7.5%. Before inserting the needle, the surface was cleaned and the first millilitre of blood was discarded in order to avoid contamination.

Organs and tissues from kids (liver, spleen, kidney, heart and skeletal muscle) were removed. Slaughter and sampling rooms were close, but distinct in order to avoid possible contamination. For the same reason, the outermost layer of tissue from each organ was removed to sample the inner part. Three aliquots (25 mg each) from each sample were immediately stored at -20°C in sterile tubes for DNA

extraction. Later on, each aliquot underwent PCR reaction. Specimens from kidney, liver, heart and muscle from right leg samples (5 g each) were washed in saline and stored at -80°C to determine enzyme activity.

DNA extraction and quantification

Milk (10 ml) and, as control, conventional and transgenic soybean meal s.e. samples (100 mg) were extracted in duplicate according to the Wizard extraction method (Promega, Madison, WI, USA). Milk samples were incubated at 4°C overnight and centrifuged ($2.000\times g$ for 20 min at 4°C) to separate cream, skimmed milk and sediment; only the sediment fraction was subjected to DNA extraction. The somatic cell pellet obtained was washed twice with phosphate buffered saline (PBS, pH 7.2). The pellet and the ground plant samples were resuspended by vortexing in 860 μl of extraction buffer [10 mM Tris-HCl (pH 8.0), 150 mM NaCl, 2 mM EDTA, 1% (w/v) SDS], 100 μl guanidine hydrochloride (5M) and 40 μl of proteinase K (20 mg/ml) and then incubated at 58°C for at least 3 h on a shaking incubator and centrifuged at $20.000\times g$ for 10 min. Five hundred microlitres of the supernatant were incubated with 5 μl RNase (10 mg/ml) at 37°C for 10 min. One millilitre of Wizard DNA purification resin (Promega) was added to the supernatant. The DNA-resin mixture was pushed through the column and washed with 2 ml 80% (v/v) isopropyl alcohol followed by centrifugation at $20.000\times g$ for 1 min. After drying at 70°C for 10 min, the DNA was eluted with 50 μl of 70°C elution buffer [10 mM Tris-HCl (pH 9.0), 0.1 mM EDTA] and centrifuged at $20.000\times g$ for 1 min.

NucleoSpin[®] Tissue and NucleoSpin[®] Blood kit (Macherey-Nagel, Duren, Germany) were used for extraction of tissue (25 mg) and blood (200 μl) samples, respectively, according to the manufacturer's protocol. Each sample was extracted in duplicate and stored at -20°C until use. In such a way, a total of six samples for each kid tissue and organ were obtained. In addition, as suggested by Nemeth (2004), negative (buffer only) control to each set of DNA extraction was included to monitor for any contamination that could have occurred during the DNA extraction procedure. The DNA was quantified by spectrophotometer (Thermo Electron Corporation, Waltham, MA, USA) according to standard molecular techniques (Sambrook *et al.*, 1989).

PCR analyses

In order to avoid contamination, PCR reactions were assembled in an ultraviolet-sterilised hood. Filter tips against sample aerosol and sterile disposable tubes were used during pipetting. All PCR amplifications were performed on a Gene Amp PCR System 2400 (Applied Biosystems, Foster City, CA, USA). DNA was amplified using Taq DNA polymerase (Invitrogen, Carlsbad, CA, USA) in accordance with the manufacturer's instructions. Sequence, amplicon size and annealing temperature of the primer pair sets (Sigma-Genosys Ltd, Haverhill, UK) used for PCR are shown in Table 2.

Primers Cap 144/496 were previously used to amplify a conserved portion of caprine mtDNA, which encodes the 12S ribosomal RNA (12S rRNA) gene of mitochondrial DNA from caprine (Bottero *et al.*, 2002).

Subsequently, samples were monitored for the presence of the chloroplast sequence for tRNA Leu by using the Clor 1/2 primers designed on the chloroplast *trnL* sequence (Terzi *et al.*, 2004). The obtained amplicon was analysed by electrophoretic separation and purified from gel. The identity was verified by sequencing and subsequently by Basic Local Alignment Search Tool (BLAST) analysis.

About the sensibility, we performed a real time PCR with Clor1/Clor2 primer and specific TaqMan probe to verify the presence of tRNA Leu gene, housekeeping gene in conventional and transgenic genotype. Then, we elaborated a standard curve and the results obtained demonstrated the efficiency of primer/probe combination to notice the presence of soybean genotype and to give quantification.

Finally, species-specific primers for conventional and GM soybean were used: Le1n02 5/3 which amplifies the soybean lectin gene (Kuribara *et al.*, 2002); 35S 1/2 and CP4 EPSPS 1/2 which amplify, respectively, part of the 35S promoter (Lipp *et al.*, 1999) derived from the cauliflower mosaic virus and part of the specific gene sequence (CP4 EPSPS) that provides herbicide tolerance derived from *Agrobacterium tumefaciens* strain CP4 both present in the genomic structure of GM soybean (Hernández *et al.*, 2003). The primer pairs were selected from those reported in the literature (Jennings *et al.*, 2003) with the aim of obtaining short amplicons (118 bp), compatible with highly fragmented DNA samples.

Table 2 Sequence (5'-3'), amplicon size (bp) and annealing temperature ($^{\circ}\text{C}$) of primer pairs used in PCR

Fragment name	Primer sequence (5'-3')	Amplicon size (bp)	T_M ($^{\circ}\text{C}$)
Cap 144	CGC CCT CCA AAT CAA TAA G	326	55
Cap 469	AGT GTA TCA GCT GCA GTA GGG TT		
Clor1	TTCCAGGGTTTCTCTGAATTTG	100	60
Clor2	TATGGCGAAATCGGTAGACG		
Le1n02-5	GCCTCTACTCCACCCCA	118	59
Le1n02-3	GCCCATCTGCAAGCCTTTT		
35S-1	GCTCCTACAAATGCCATCA	195	56
35S-2	GATAGTGGGATTGTGCGTCA		
CP4EPSPS 1	GCA AAT CCT CTG GCC TTT CC	145	60
CP4EPSPS 2	CTT GCC CGT ATT GAT GAC GTC		

A 100 base pairs (bp) ladder, containing linear DNA fragments, served as size standard reference. The PCR was done three times, and samples with positive results at least twice were judged as positive (Chowdhury *et al.*, 2003b). In every PCR run, positive and negative controls were included to ensure reproducibility and absence of contaminants. For positive controls, reference DNA consisting of purified conventional and transgenic soybean DNA was amplified in parallel with the samples to ensure correct performance of the PCR; for negative control (buffer blank), water instead of DNA was added to the PCR mix to check for cross contamination with soybean DNA in the PCR mix or its constituents (Klaften *et al.*, 2004).

Enzyme assay

The levels of aspartate aminotransferase (AST), alanine aminotransferase (ALT), creatine kinase, lactic dehydrogenase (LDH), gamma glutamyltransferase (GGT) and alkaline phosphatase were determined in serum and homogenates prepared from samples of liver, kidney, heart and skeletal muscle. Briefly, tissue (1 g) was homogenised in ice-cold homogenisation buffer (in mM): 280 mannitol, 10 KCl, 1 MgCl₂, 0.2 Pefabloc SC, 10 Hepes, pH 7.0 adjusted to pH 7 with Tris HCl 10 mM. After centrifugation at 10 000×g for 10 min the upper layer was used for analysis.

Enzyme activity was determined spectrophotometrically (340 nm) by using reagents from Spinreact SA (Sant Esteve de Bas, Spain). Determinations were performed according to the manufacturer's instructions, including the conversion of the results into units/litre. In order to assess the isoenzymatic distribution of LDH, electrophoretic separation was performed on each sample. Briefly, 20 µl of sample were applied on cellulose acetate membranes and electrophoresis was performed under denaturing conditions at 200 V for 50 min in barbital buffer. The substrate (tris pH 7.4, sodium lactate, sodium azide) was incorporated in agarose and put on the acetate membrane immediately after electrophoresis. Therefore, it was possible to observe LDH isoenzymes (ISO-LAD commercial kits by Chemetron Chimica S.p.A., Milan, Italy). Percentage of isoenzyme fractions of total LDH were quantified by using a densitometer (CGA, Florence, Italy). Data were then converted into units/litre (serum) or units/gram of tissue (organs).

LDH histochemistry

Samples were prepared using a cryostat: 10 µm sections were obtained, with five serial sections put in every slide. The frozen sections were air-dried and then incubated, always in the dark, in LDH-incubating medium at 37°C for 10 min. The incubated medium consisted of 50 mM Tris-HCL (pH 7.4) containing 67 mM sodium lactate (L 1375, Sigma-Aldrich, Milan, Italy), 5 mM nitroblue tetrazolium (N 6876, Sigma), 5 mM MgCl₂ and 3 mM NAD (N1511, Sigma). Each sample, both from control and treated animals, was compared with its negative control section, which was obtained by incubation in incubating medium without sodium lactate. The histochemical reaction was stopped, at the same time for every slide, by fixing the section in formalin for 15 min at room temperature. The sections were then counterstained with methyl green. The slides

were mounted in PBS/glycerol (1:1) and then examined and photographed. Five slides for each examined tissue were independently evaluated by two observers using a DMRA2 microscope (Leica Microsystems Wetzlar GmbH, Germany).

Statistics

Differences in kid slaughter weight as well as spleen, liver, kidney and heart were tested by one way analysis of covariance (control *v.* treated group). Data were covariates for birth weight to reduce variations from the mean square error in the dependent variables. For goat milk and blood, differences in gene detection among sampling periods were tested by the Cochran-Q test. The presence of plant DNA fragments in kids blood, skeletal muscle and organs was analysed by using the χ^2 test. Results for enzyme assays were expressed as mean \pm standard deviation and differences between groups were analysed by the Student's *t*-test. All statistics were performed with SPSS 12.1 software (SPSS, 1999).

Results

BW of the kids at birth was on average 4.2 *v.* 3.7 kg, respectively for groups C and T. As a consequence, birth weight was used to covariate slaughter and organ weights. Neither body nor organ weights significantly differed between groups (Table 3).

Detection of DNA fragments

The quality of each DNA sample extracted from milk and blood of control and treated goats and from liver, kidney, skeletal muscle, spleen, heart and blood of each group of kids was first verified using the Cap 144/496 primers that were used to amplify a conserved portion of caprine mtDNA 12S rRNA sequence. An example is reported in Figure 1.

Table 3 Body and organ weights in control (n = 10) and treated (n = 10) kids fed RoundUp Ready[®] soybean meal

	Control	Treated	m.s.e.
BW (kg)	11.68	10.73	1.64
Spleen (g)	36.03	30.83	42.81
Liver (g)	312.2	243.3	1898.6
Kidney (g)	77.46	59.11	167.3
Heart (g)	106.4	69.61	1285.1

All data were collected at slaughter.




Figure 1 Representative electrophoresis gels of amplified DNA in: (left) milk and blood from control (lines 1 and 3) and treated (lines 2 and 4) goats; (right) liver, kidney, skeletal muscle, spleen, heart and blood from control (lines 1 to 6, respectively) and treated (lines 7 to 12, respectively) kids. In each panel, lane M contains a 100 bp DNA ladder; '-' is a negative control (no DNA template).

Table 4 Number of animals in which DNA sequences were detected. Blood and milk derived from dams which either (treated, $n = 10$) or not (control, $n = 10$) were fed Roundup Ready soybean meal s.e.

	Blood										Milk										Significance
	Control					Treated					Control					Treated					
	−60	−30	0	30	60	−60	−30	0	30	60	15	30	45	60	75	15	30	45	60	75	
Day	−60	−30	0	30	60	−60	−30	0	30	60	15	30	45	60	75	15	30	45	60	75	
Chlor	10	8	10	6	8	10	9	10	9	9	7	8	8	8	8	8	9	10	8	8	**
Lectine	−	5	5	5	5	−	5	5	5	5	5	6	6	7	7	5	6	6	7	7	**
35S	−	−	−	−	−	−	2	3	3	5	−	−	−	−	−	3	4	6	6	7	**
CP4 <i>EPSPS</i>	−	−	−	−	−	−	2	4	5	7	−	−	−	−	−	4	5	5	5	5	**

chlor = chloroplast DNA fragment.

** $P < 0.01$.**Table 5** Number of dams producing milk at 60 days in lactation in which DNA sequences were detected. Number of kids with organs in which DNA sequences were detected

	Control							Treated						
	Milk		Kids					Milk		Kids				
	Day 60	Blood	Muscle	Kidney	Liver	Spleen	Heart	Day 60	Blood	Muscle	Kidney	Liver	Spleen	Heart
Chlor	8	7**	7**	8**	8**	7**	6**	8	7**	8**	8**	6**	7**	7**
Lectine	7	4*	4*	5**	5**	3	3	7	4*	4*	5**	5**	2	4*
35S	—	—	—	—	—	—	—	6	4**	3	5**	5**	2	3
CP4 EPSPS	—	—	—	—	—	—	—	5	2	4**	3*	3*	2	4**

Milk derived from dams which either (treated, $n = 10$) or not (control, $n = 10$) were fed Roundup Ready soybean meal s.e. Organs derived from their nursing kids. The lectine, 35S and CP4 EPSPS fragments were investigated only in those samples, which were positive for the chloroplast DNA fragment (Chlor).* $P < 0.05$; ** $P < 0.01$. This P -value indicates a significant proportion of the kids found positive for the same DNA fragments detected in milk.

Table 4 shows the number of goats in which different DNA sequences were detected. PCR was done three times for each sample and only samples with at least twice a positive result were judged as positive (Chowdhury *et al.*, 2003b). A chloroplast specific DNA fragment (*trnL*, 100 bp) was found, both in blood and milk, in the majority of control and treated samples. In the samples, which were positive for the presence of chloroplast DNA, the fragments of the lectin gene, 35S promoter and CP4 EPSPS were investigated and, again, several control and treated samples were found positive for lectin gene fragments. Transgenic target DNA sequences (35S and CP4 EPSPS) were not detected in blood and milk from control goats that received a diet containing conventional soybean meal. By contrast, transgenic DNA fragments were amplified from samples (blood and milk) from goats that received transgenic soybean. The Cochran's Q-test revealed that detection of chloroplast, lectin, 35S promoter and CP4 EPSPS gene fragments in goats' milk and blood was significant ($P < 0.01$) during the whole experiment.

Table 5 shows the number of kids from control and treated group in which different DNA sequences were detected. The chloroplast specific DNA fragment (*trnL*, 100 bp) was found, in all organs, in the majority of control and treated samples. In the samples, which proved positive for the presence of chloroplast DNA, the fragments of lectin gene, 35S promoter and CP4 EPSPS were investigated and, again, several control and treated samples were found positive for lectin gene fragments. Transgenic target DNA sequences (35S and CP4 EPSPS) were not detected in organs from control kids whose

mothers received a diet containing conventional soybean meal. In contrast, transgenic DNA fragments were amplified from some samples of kids whose mothers received transgenic soybean. As seen, the CP4 EPSPS fragment was found in more samples compared with the 35S fragment, the reason for such result is not clear, but it could be due to slight differences in PCR conditions. In order to assess the significance of the presence of transgenic DNA fragments in kids organs, a χ^2 analysis of all kids data v. those detected in milk from the corresponding mothers at day 60 (slaughtering time) was performed and revealed that: (i) chloroplastid fragment presence was significant in all organ and tissues from both control and treated groups, (ii) lectin fragments detection was significant in kidney and liver of both groups ($P < 0.01$), in blood and muscle from control group ($P < 0.05$) and in blood, muscle and heart from treated group ($P < 0.05$), (iii) the detection of 35S promoter was significant ($P < 0.01$) in liver, kidney and blood of treated kids and (iv) the detection of CP4 EPSPS gene fragment was significant in liver and kidney ($P < 0.05$) and in heart and muscle ($P < 0.01$) from treated group.

In Figure 2, a representative example is presented illustrating the detection of endogenous high copy chloroplast specific DNA fragment (*trnL*, 100 bp) in goats milk and blood of both groups and in tissue, organs and blood from both control and treated groups of kids.

In Figure 3, a representative example obtained by using Le1n02 5/3 soybean specific primers shows the lectin signal detected in goats blood and milk samples and in the plant control, as well as in control and treated groups of kids.




Figure 2 Confirmation of amplified endogenous chloroplast specific DNA fragments, trnL (representative data) in: (left) milk and blood from control (lines 1 and 3) and treated (lines 2 and 4) goats; (right) liver, kidney, skeletal muscle, spleen, heart and blood from control (lines 1 to 6, respectively) and treated (lines 7 to 12, respectively) kids. In each panel, lane M contains a 100 bp DNA ladder; '–' is a negative control (no DNA template) and '+' is a positive control (DNA extracted from soybean meal).



Figure 3 Representative electrophoresis analysis of single copy specific soybean DNA fragment, lectin gene in: (left) milk and blood from control (lines 1 and 3) and treated (lines 2 and 4) goats; (right) liver, kidney, skeletal muscle, spleen, heart and blood from control (lines 1 to 6, respectively) and treated (lines 7 to 12, respectively) kids. In each panel, lane M contains a 100 bp DNA ladder; '–' is a negative control (no DNA template), and '+' is a positive control (DNA extracted from soybean meal).




Figure 4 Representative data of amplified transgenic DNA fragments. (a) 35S promoter fragments (195 bp) in milk and blood from control (lines 1 and 2, respectively) and treated (lines 3 and 4, respectively) goats. (b) 35S promoter fragments (195 bp) in liver, kidney, spleen, heart, skeletal muscle and blood from control (lines 7 to 12, respectively) and treated (lines 1 to 6, respectively) kids. (c) fragments of CP4 EPSPS gene (145 bp) in milk and blood from control (lines 3 and 4, respectively) and treated (lines 1 and 2, respectively) goats. (d) fragments of CP4 EPSPS gene (145 bp) in liver, kidney, spleen, heart, skeletal muscle and blood from control (lines 1 to 6, respectively) and treated (lines 7 to 12, respectively) kids. In each panel, lane M contains a 100-bp DNA ladder, '–' is a negative control (no DNA template), and '+' is a positive control (DNA extracted from Roundup Ready soybean meal).

Transgenic target DNA sequences (35S and CP4 EPSPS) detected in either goats that received a diet containing transgenic soybean meal or their kids as well as in the RoundUp Ready soybean positive control are represented in Figure 4. In particular, in blood and milk of treated goats, Figure 4 shows fragments of 35S promoter and fragments of the CP4 EPSPS gene. In treated kids, a representative detection of fragments of 35S and the CP4 EPSPS gene is reported.

Since it could be likely that the DNA detected may be too small to be of any biological significance, we later

investigated whether larger DNA fragments could be detected in milk samples. To this purpose, a primer amplifying a fragment of 447 bp of the p-35S/CP4 EPSPS gene (Cardarelli *et al.*, 2005) was used. In 57% of the samples that were found positive for smaller transgenic fragment, such sequence was detected (see addendum).

Enzyme activity

As depicted in Table 6, AST and ALT were significantly lower in serum from goats fed GM soybean at day 60 and at days 30 and 60, respectively. Overall, very limited differences in the monitored enzyme activity have been observed.

Figure 5 shows the differences in enzyme activity in kids' heart, skeletal muscle, kidney, liver and serum. Statistical differences ($P < 0.05$) were detected in kidney for GGT and LDH, whereas in the heart and skeletal muscle this result was seen only for LDH ($P < 0.05$). Hence, in five of the 30 cases presented, differences were observed.

No statistical differences were found for studied enzymes in serum and liver. The increase in LDH activity was confirmed by histochemistry (Figure 6). LDH histochemistry showed a widespread distribution of enzyme activity in all the tissues that were examined. The expression of LDH activity was further illustrated by a violet staining in the cytoplasm of the cells. In tissue sections used as negative controls this staining was light blue. Myocytes, myocardiocytes, epithelial cells of renal tubules and hepatocytes all expressed LDH activity. LDH staining was more intense in myocytes, myocardiocytes and epithelial cells of renal tubules of treated than in those of control animals. Arrows show some zones where LDH histochemical staining is more intense. No difference was observed between the two groups of animals in hepatocytes.

Table 7 shows the relative distribution of LDH isoenzymes in serum and in tissues. Significant differences ($P < 0.05$) between control and treated animals were detected for heart and kidney LDH1 and for muscles LDH1 and LDH2, thus confirming the significant increase of the enzyme in these tissues.

In addition, analysis of isoenzyme distribution revealed that LDH1 significantly increased also in the liver if we compare the proportional distribution of isoenzymes (51.2 ± 3.7 v. 57.8 ± 1.4). Moreover, heart LDH2 (4.4 ± 1.9 v. 2.9 ± 0.9) and LDH3 (1.9 ± 0.4 v. 1.1 ± 0.7) significantly decreased and a similar result was seen for muscle LDH5 (55.6 ± 5.7 v. 49.4 ± 2.8).

Discussion

In our study, the weight of the organs of kids did not differ between groups, in accordance to findings of previous research (Tudisco *et al.*, 2006b) carried out on rabbits fed GM soybean meal.

Detection of DNA fragments

As stated by the EFSA (2008), several aspects have to be investigated when considering whether or not recombinant DNA from GM plants, or the derived proteins can end up in

Table 6 Monthly levels of AST, ALT, CK, LDH, GGT and ALP in serum taken 60 days before to 60 days after lambing from control (C) and treated (T) group of goats

Enzyme (U/l)	Group	Day									
		-60		-30		0		30		60	
		Mean	s.d.	Mean	s.d.	Mean	s.d.	Mean	s.d.	Mean	s.d.
CK	C	64.4	15.77	59.7	15.38	59.5	14.89	58.1	15.52	59.3	19.71
	T	58.5	19.36	60.0	15.51	57.2	16.67	58.1	15.19	53.7	15.64
AST	C	137.7	17.59	143	30.1	137.2	27.71	125.5	30.18	146	36.9
	T	138.9	37.79	135.1	34.21	138.1	26.63	115.5	20.61	121.5*	22.56
ALT	C	22	10.3	24.1	9.57	21.8	7.00	27.4	7.12	24.1	7.32
	T	21.4	2.49	20.5	6.32	19.9	4.67	20.5*	8.17	19.3*	4.00
ALP	C	301.5	69.36	286.5	112.28	265.5	90.23	335	115.7	294.6	95.64
	T	289.1	98.88	238.8	63.42	286.7	73.12	253.5	68.32	216.8	93.00
GGT	C	40.7	13.28	47.8	26.52	43.6	18.94	40.6	21.58	39.4	15.18
	T	37.8	13.47	51	12.3	41	18.7	46.7	14.56	42.9	19.88
LDH	C	1034.7	373.38	1072	452.0	1059	423.6	969	460.0	1065.2	374.63
	T	1028.6	385.00	1014.9	316.11	1103.1	554.71	963.4	494.08	1013.7	301.00

AST = aspartate aminotransferase; ALT = alanine aminotransferase; CK = creatine kinase; LDH = lactic dehydrogenase; GGT = gamma glutamyltransferase; ALP = alkaline phosphatase.

* $P < 0.05$ control (C) v. treated (T) dams.




Figure 5 Levels of creatine kinase (CK), lactic dehydrogenase (LDH), aspartate aminotransferase (AST), alanine aminotransferase (ALT), gamma glutamyltransferase (GGT) and alkaline phosphatase (ALP) in serum and in homogenates from liver, kidney, heart and skeletal muscle from control (▨) and treated (■) kids. * $P < 0.05$.

animal tissues and products. These include (i) the fate of the recombinant DNA and protein during feed processing and ensiling; (ii) the fate of the recombinant DNA and protein

in the gastrointestinal tract of animals fed with GM feed; (iii) the potential absorption of the digested pieces of DNA or protein into animal tissues/products and (iv) the potential




Figure 6 LDH histochemistry on negative control sections (a, d, g, l), control (b, e, h, m) and treated (c, f, i, n) animals. LDH activity was revealed by a violet staining. In negative control sections, a light bluish staining was observed. In the skeletal muscle, heart and epithelial cells of the renal tubule of treated animal (c, f, i) the intensity of LDH staining (arrows) was higher than in the same tissues (b, e, h) of control animals. Bar = 50 μ m.

Table 7 Relative distribution of LDH isoenzymes in serum (S) and in homogenates from heart (H), SM, kidney (K) and liver (L) from control (C) and treated (T) group of kids

Tissue	Group	Relative distribution of LDH isoenzymes (U/g)											
		Total LDH activity		LDH-1		LDH-2		LDH-3		LDH-4		LDH-5	
		Mean	s.d.	Mean	s.d.	Mean	s.d.	Mean	s.d.	Mean	s.d.	Mean	s.d.
S (U/L)	C	913	148.2	23.2	6.89	47.6	11.00	56.4	7.36	209.9	31.00	575.8	103.00
	T	873	134.7	24.2	7.56	46.3	9.11	55.6	15.42	209.4	27.00	537.5	87.46
H (U/g)	C	245.9	63.77	224.5	57.82	11	3.8	4.6	1.00	2.9	3.61	2.8	2.71
	T	318.2*	57.39	300.2*	52.71	9	2.4	3.8	2.87	2.5	1.47	2.6	1.38
SM (U/g)	C	403	76.71	56.2	10.19	45.9	36.12	36.1	8.78	39	13.0	226.5	59.00
	T	517*	132.52	100.5*	35.61	61.2*	17.33	51	23.1	48.8	18.00	255.5	67.81
K (U/g)	C	47	9.0	27.8	5.78	11.8	2.67	4.9	1.13	1.3	0.51	1.2	0.74
	T	66.8*	16.53	42.5*	10.89	15.5	4.48	6.1	2.11	1.4	0.78	1.3	0.52
L (U/g)	C	134.7	36.18	68.7	17.42	25.5	6.21	16.7	6.56	12.2	5.11	11.7	6.31
	T	144.2	27.00	83.5	17.46	26.8	1.72	16.2	4.00	8.8	3.33	8.9	4.56

LDH = lactic dehydrogenase; SM = skeletal muscle.

* $P < 0.05$ control (C) v. treated (T) dams.

of biological functionality of absorbed DNA and protein fragments.

Our findings show plant DNA fragments are likely to survive digestive processes to some extent (Duggan *et al.*, 2003; Einspanier *et al.*, 2004), as well as their transfer to blood and

milk. In addition, the detection of plant DNA in tissues and organs of nursed kids could support the hypothesis of a gene transfer through milk. Chloroplast target DNA sequences were found in our previous research in blood and tissues of lambs allowed to nurse from the dam. However, no single

copy feed fragment was detectable in lamb samples nor in the mother's milk and blood. The reason for this high recovery of chloroplast DNA in animal tissues is a consequence of the larger quantities of these genes as compared with transgenic genes, which are usually found as a single copy gene (Aumaitre *et al.*, 2002).

Although few have found recombinant DNA sequences in any organ or tissue sample from animals fed GM plants (EFSA, 2007), our results showed that small DNA fragments can also be detected in kids organs when mothers are fed GM soybean. Recombinant plant DNA has also been found by other authors as a rare event (Chowdhury *et al.*, 2003a; Mazza *et al.*, 2005; Sharma *et al.*, 2006) in organs and tissues of animals fed GM feeds. On the contrary, studies aiming at verifying the presence of transgenic fragments in milk gave negative results. In dairy cows fed GM soybean the chloroplastid sequence was found in leucocytes but not in milk (Klotz and Einspanier, 1998). In other trials carried out with dairy cows fed GM soybean meal and maize (Phipps *et al.*, 2003; Nemeth *et al.*, 2004) or goats fed GM maize (Rizzi *et al.*, 2008), the chloroplast DNA fragments were detected in milk, but no transgenic fragments were found.

Detection of GM sequence in animal organs and tissues alone cannot justify the public concern regarding the human consumption of products from farm animals fed transgenic crops. Indeed, the implications of DNA transfer from food containing GM plants to an organism is no higher as compared with DNA transfer from food containing the corresponding conventional plant (Mazza *et al.*, 2005).

Enzyme assay

In goats, a significant decrease in AST (day 60) and ALT (days 30 and 60) was detected in serum from goats fed GM soybean, but in both groups enzyme levels were in the normal range. Therefore, such difference should have no clinical relevance. Similarly, lower levels of ALT were found in serum from buffaloes fed transgenic cottonseed and the authors concluded that there were no differences in buffalo health status as determined by haematobiochemical constituents (Singh *et al.*, 2003). On all other days, and also in kids, the levels of the enzyme tested in serum did not show significant differences, thus suggesting that no adverse effects were induced by GM feed in treated animals, as previously reported in dairy cows (Yonemochi *et al.*, 2003). By contrast, analysis of enzyme total and relative activities in tissues gave a different picture. As depicted in Figure 5 significant differences in enzyme levels concerned the kidney (higher levels of LDH and GGT) as well as heart and skeletal muscle (higher levels of LDH) in kids drinking milk from treated animals. This seems to indicate that some alteration occurred in kids even if serum levels were not affected. Moreover, LDH significantly increased in three organs (kidney, heart and skeletal muscle), suggesting that enzyme synthesis is generally altered in kids fed milk from GM-fed dams. The relative distribution of LDH isoenzymes confirms this hypothesis, showing significant differences in heart and kidney LDH1 and in muscle LDH1 and LDH2. In addition, if we compare

the distribution of isoenzymes in percentage terms, LDH1 significantly increased also in the liver and heart LDH2 and LDH3 significantly decreased; a similar result was also seen for muscle LDH5. Therefore, the significant increase in LDH1 was accompanied by a percent decrease of other isoenzymes thus suggesting that, since LDH is a tetrameric enzyme made up of M and H subunits, a different combination of such subunits occurred in the heart (LDH-H₄ > LDH-H₃M₁ and LDH-H₂M₂) and in the skeletal muscle (LDH-H₄ and LDH₃M₁ > LDH-M₄). A similar shift has been previously described in rabbits (LDH-H₄ > LDH-H₁M₃) where the higher amount of H subunit was assumed to mean a higher specificity for the reduction of alpha-hydroxybutyrate to alpha-oxobutyrate, thereby showing that some metabolic changes occurred in the liver (Tudisco *et al.*, 2006b). Our data confirm this hypothesis: the isoenzyme mainly composed by the H subunit showed an increased percentage, which was also found in the muscle where LDH1 is not the dominant isoenzyme. Therefore, increased activity of LDH1 occurred in organs of GM-fed kids, as confirmed by histochemistry which showed a higher expression of total LDH in heart, kidney and skeletal muscles.

Since LDH1 is known to be involved in cell metabolism by favouring the reaction of lactate to pyruvate (Van Hall, 2000), our results could indicate a general increase in cell metabolism. This hypothesis is in agreement not only with results obtained in rabbits but also with those who found significant modifications of some nuclear features in GM-fed mice suggesting a high metabolic rate and intense molecular trafficking (Malatesta *et al.*, 2002). In any event, there is no scientific evidence that a local increase in LDH metabolism may be dangerous for health especially when it does not affect serum levels. Moreover, since serum activities of all the enzymes showed similar levels between the groups, it would be over speculative to assert that the GM diet was responsible for a local increase in LDH metabolism. However, the synthesis of LDH did actually change in more than one organ and such results should be taken into account for future research.

The mere detection of recombinant DNA fragments in animal organs and tissues cannot justify, by itself, public concerns regarding human consumption of products from farm animals fed transgenic crops. By contrast, although both BW and organ weight did not significantly differ between groups, any alteration in cell metabolism should be taken into account in this field. The modification in LDH synthesis, even without affecting serum levels of the enzyme, suggests an increase in cell metabolism. Therefore, possible long-term effects of such an alteration need to be elucidated.

In conclusion, our research attempted a wider survey in terms of detection of DNA fragments as well as investigating the impact of GM feed in terms of health. Our results showed that small DNA fragments can be detected in milk but also in kids organs when mothers are fed GM soybean and revealed elevated levels of LDH in tissues, suggesting an increase in cell metabolism. Although our data seem to conflict with the majority of studies in this field, they are consistent with other studies and the longer term consequences to health following GM food intake merit further consideration.

References

- AOAC 2000. Official methods of analysis, 17th Edition. Association of Official Analytical Chemists, Arlington, Virginia.
- Aumaitre A, Aulrich K, Chesson A, Flachowsky G and Piva G 2002. New feeds from genetically modified plants: substantial equivalence, nutritional equivalence, digestibility, and safety for animals and the food chain. *Livestock Production Science* 74, 223–238.
- Bottero MT, Civera T, Numera D, Rosati S, Sacchi P and Turi RM 2002. A multiplex polymerase chain reaction for the identification of cow's, goat's and sheep's milk in dairy products. *International Dairy Journal* 13, 277–282.
- Cardarelli P, Branquinho MR, Ferreira RTB, da Cruz FP and Gemal AL 2005. Detection of GMO in food products in Brazil: the INCQS experience. *Food Control* 16, 859–866.
- Chowdhury EH, Kuribara H, Hino A, Sultana P, Mikami O, Shimada N, Guruge KS, Saito M and Nakajima Y 2003a. Detection of corn intrinsic and recombinant DNA fragments and Cry1Ab protein in the gastrointestinal contents of pigs fed genetically modified corn Bt11. *Journal of Animal Science* 81, 2546–2551.
- Chowdhury EH, Mikami O, Nakajima Y, Kuribara H, Hino A, Suga K, Hanazumi M and Yomemochi C 2003b. Detection of genetically modified maize DNA fragments in the intestinal contents of pigs fed StarLink™ CBH351. *Veterinary and Human Toxicology* 45, 95–96.
- Duggan PS, Chambers PA, Heritage J and Forbes JM 2003. Fate of genetically modified maize DNA in the oral cavity and rumen of sheep. *British Journal of Nutrition* 89, 159–166.
- EFSA 2007. Statement on the fate of recombinant DNA or proteins in meat, milk and eggs from animals fed with GM feed. Retrieved July 19, 2007, from http://efsa.europa.eu/EFSA/efsa_locale
- EFSA GMO Panel Working Group on Animal Feeding Trials 2008. Safety and nutritional assessment of GM plants and derived food and feed: the role of animal feeding trials. *Food and Chemical Toxicology* 46 (suppl. 1), S2–S70.
- Einspanier R, Lutz B, Rief S, Berezina O, Zverlov V, Schwarz W and Mayer J 2004. Tracing residual recombinant feed molecules during digestion and rumen bacterial diversity in cattle fed transgene maize. *European Food Research and Technology* 218, 269–273.
- Hernández M, Rodríguez-Lázaro D, Esteve T, Prat S and Pla M 2003. Development of melting temperature-based SYBR Green I polymerase chain reaction methods for multiplex genetically modified organism detection. *Analytical Biochemistry* 323, 164–170.
- INRA 1978. Alimentation des ruminants. Edition INRA, Paris, France.
- Jennings JC, Kolwyck DC, Kats SB, Whetsell AJ, Surber JB, Cromwell GL, Lirette RP and Glenn KC 2003. Determining whether transgenic and endogenous plant DNA and transgenic protein are detectable in muscle from swine fed Roundup Ready soybean meal. *Journal of Animal Science* 81, 1447–1455.
- Klaften M, Whetsell A, Webser J, Grewal R, Fedyk E, Einspanier R, Jennings J, Lirette R and Glenn K 2004. Animal biotechnology: challenges and prospects. In ACS symposium series (ed. MM Bhalgat, WP Ridley, AS Felsot and JN Seiber), pp. 83–99. American Chemical Society, Washington, DC.
- Klotz A and Einspanier R 1998. Detection of "novel-feed" in animals? Injury of consumers of meat and milk is not expected. *Mais* 3, 109–111.
- Kuribara H, Shindo Y, Matsuoka T, Futo S, Aoki N, Hirao T, Ariyama H, Goda Y, Toyoda M and Hino A 2002. Novel reference molecules for quantitation of genetically modified maize and soybean. *Journal of AOAC International* 85, 1077–1089.
- Laudadio V, Petrerà F, Tudisco R and Infascelli F 2006. Genetically modified soya bean in sheep feeding: detection of DNA fragments in suckling lambs. *Proceedings of 12th AAAP Animal Science Congress*, 18–22 September, Bexco, Busan, Korea, pp. 3–6.
- Lipp M, Brodmann P, Pietsch K, Pauwels J and Anklam E 1999. IUPAC: collaborative study of a method to detect genetically modified soybean and maize in dried powder. *Journal of AOAC International* 82, 923–928.
- Malatesta M, Caporaloni C, Gavaudan S, Rocchi MB, Serafini S, Tiberi C and Gazzanelli G 2002. Ultrastructural morphometrical and immunocytochemical analyses of hepatocyte nuclei from mice fed on genetically modified soybean. *Cell Structure and Function* 27, 173–180.
- Malatesta M, Boraldi F, Annovi G, Baldelli B, Battistelli S, Biggiogera M and Quaglino D 2008. A long-term study on female mice fed on a genetically modified soybean: effects on liver ageing. *Histochemistry and Cell Biology* 130, 967–977.
- Mazza R, Soave M, Morlacchini M, Piva G and Marocco A 2005. Assessing the transfer of genetically modified DNA from feed to animal tissues. *Transgenic Research* 14, 775–784.
- Nemeth A, Wurz A, Artim L, Charlton S, Dana G, Glenn K, Hunst P, Jennings J, Shilito R and Song P 2004. Sensitive PCR analysis of animal tissue samples for fragments of endogenous and transgenic plant DNA. *Journal of Agricultural and Food Chemistry* 52, 6129–6135.
- Padgett SR, Kolacz KH, Delannay X, Re DB, LaVallee BJ, Tinus CN, Rhodes WK, Otero YI, Barry GF, Eichholtz DA, Peschke VM, Nida DL, Taylor NB and Kishore GM 1995. Development, identification and characterization of a glyphosate-tolerant soybean line. *Crop Science* 35, 1451–1461.
- Phipps RH, Deaville ER and Maddison BC 2003. Detection of transgenic and endogenous plant DNA in rumen fluid, duodenal digesta, milk, blood, and feces of lactating dairy cows. *Journal of Dairy Science* 86, 4070–4078.
- Reuter T and Aulrich K 2003. Investigations on genetically modified maize (Bt-maize) in pig nutrition: fate of feed-ingested foreign DNA in pig bodies. *European Food Research and Technology* 216, 185–192.
- Rizzi A, Brusetti L, Arioli S, Nielsen KM, Tamagnini I, Tamburini A, Sorlini C and Daffonchio D 2008. Detection of feed-derived maize DNA in goat milk and evaluation of the potential of horizontal transfer to bacteria. *European Food Research and Technology* 227, 1699–1709.
- Sambrook J, Fritsch EF and Maniatis T 1989. Molecular cloning. A laboratory manual. Cold Spring Harbor Laboratory Press, Cold Spring Harbor.
- Sharma R, Damgaard D, Alexander TW, Dugan MER, Aalhus JL, Stanford K and McAllister TA 2006. Detection of transgenic and endogenous plant DNA in digesta and tissues of sheep and pigs fed Roundup Ready canola meal. *Journal of Agricultural Food Chemistry* 54, 1699–1709.
- Singh M, Tiwari DP, Kumar A and Ravi Kumar M 2003. Effect of feeding transgenic cottonseed vis-a-vis non-transgenic cottonseed on haematobiochemical constituents in lactating Murrah buffaloes. *Asian-Australian Journal of Animal Science* 16, 1732–1737.
- SPSS Inc 1999. SPSS for windows, ver. 12.0. SPSS Inc., Chicago, IL, USA.
- Terzi V, Infascelli F, Tudisco R, Russo G, Stanca AM and Faccioli P 2004. Quantitative detection of Secale cereale by real-time PCR amplification. *LWT-Food Science and Technology* 37, 239–246.
- Tony MA, Butschke A, Broll H, Grohmann L, Zagon J, Halle I, Dänicke S, Schauzu M, Hafez HM and Flachowsky G 2003. Safety assessment of Bt 176 maize in broiler nutrition: degradation of maize-DNA and its metabolic fate. *Archives of Animal Nutrition* 57, 235–252.
- Tudisco R, Infascelli F, Cutrignelli MI, Bovera F, Morcia C, Faccioli P and Terzi V 2006a. Fate of feed plant DNA monitored in water buffalo (*Bubalus bubalis*) and rabbit (*Oryctolagus cuniculus*). *Livestock Science* 105, 12–18.
- Tudisco R, Lombardi P, Bovera F, D'Angelo D, Cutrignelli MI, Mastellone V, Terzi V, Avallone L and Infascelli F 2006b. Genetically modified soybean in rabbit feeding: detection of DNA fragments and evaluation of metabolic effects by enzymatic analysis. *Animal Science* 82, 193–199.
- Van Hall G 2000. Lactate as a fuel for mitochondrial respiration. *Acta Physiologica Scandinavica* 168, 643–656.
- Van Soest PJ, Robertson JB and Lewis BA 1991. Methods for dietary fiber, neutral detergent fiber, and nonstarch polysaccharides in relation to animal nutrition. *Journal of Dairy Science* 74, 3583–3598.
- Yonemochi C, Ikeda T, Harada C, Kusama T and Hanazumi M 2003. Influence of transgenic corn (CBH 351, named Starlink) on health condition of dairy cows and transfer of Cry9C protein and cry9C gene to milk, blood, liver and muscle. *Animal Science Journal* 74, 81–88.